

AFRL-RW-EG-TP-2008-7420

CRYSTAL FIELD INTERACTIONS OF “LARGE” ROTORS EMBEDDED IN SOLID PARAHYDROGEN (BRIEFING CHARTS AND ABSTRACT)

C. Michael Lindsay
Mario Fajardo
Air Force Research Laboratory
Munitions Directorate
AFRL/RWME
Eglin AFB, FL 32542-6810



SEPTEMBER 2008

BRIEFING CHARTS AND ABSTRACT

The briefing charts were presented at the 63rd Ohio State University International Symposium on Molecular Spectroscopy held 16-20 June 2008, in Columbus, Ohio. The abstract shown in block 14 of attached Report Documentation Page is published in the conference proceedings found online in the Ohio State University's Knowledge Bank, <http://hdl.handle.net/1811/33333>.

The authors are U.S. Government employees working within the scope of their positions. Therefore, this is a work of the U.S. Government and is not subject to copyright in the United States.

This paper is published in the interest of the scientific and technical information exchange. Publication of this paper does not constitute approval or disapproval of the ideas or findings.

DISTRIBUTION A: Approved for public release; distribution unlimited.
Briefing Charts approval confirmation 96 ABW/PA # 06-05-08-287; dated 5 June 2008.
Abstract approval confirmation AAC/PA # 03-05-07-141; dated 5 March 2007.

AIR FORCE RESEARCH LABORATORY, MUNITIONS DIRECTORATE

Air Force Materiel Command

■ United States Air Force

■ Eglin Air Force Base

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)	2. REPORT TYPE	3. DATES COVERED (From - To)		
4. TITLE AND SUBTITLE		5a. CONTRACT NUMBER		
		5b. GRANT NUMBER		
		5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)		5d. PROJECT NUMBER		
		5e. TASK NUMBER		
		5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)	
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT				
13. SUPPLEMENTARY NOTES				
14. ABSTRACT				
15. SUBJECT TERMS				
16. SECURITY CLASSIFICATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON 19b. TELEPHONE NUMBER (Include area code)
a. REPORT	b. ABSTRACT	c. THIS PAGE		



Crystal Field Interactions of “Large” Rotors Embedded in Solid Parahydrogen

C. Michael Lindsay, Ph.D.

Mario Fajardo, Ph.D.

Energetic Materials Branch

Munitions Directorate

Air Force Research Laboratory

DISTRIBUTION A: Approved for public release;
distribution unlimited. 96 ABW/PA 06-05-08-287




Rotation in the Solid Matrices

- Rare gas matrices and traditional molecular matrices:
 - Small hydrides: HX, H₂O, NH₂, NH₃, CH₃, CH₄
 - *Perhaps* CIF, CN (ESR and IR spectra conflict)
 - fcc lattice structure
- Solid Parahydrogen:
 - Very low angular anisotropy, and weak intermolecular interactions
 - Small hydrides: HX, H₂O, NH₃, CH₃, CH₄
 - “Large” molecules: N₂, O₂, CO, HCN, ... more?
 - fcc (as deposited) and hcp (annealed) lattice structure



Narrow spectral lines widths and the ability to examine the rotation of larger molecules allows us to examine the impurity-matrix interactions with remarkable clarity

Rapid vapor deposition of p-H₂

- Reflection based set-up (FTIR).
- [o-H₂] < 100 ppm.
- p-H₂ deposition rate ~150 mmol/hr → ~1 mm/hr thickness.
- Deposition at 2.3 K. Sample annealed at 4.3 K for 30 min.





Polarization Spectroscopy



Polarization analysis provides:

- An assignment of the perpendicular and parallel components to the crystal field perturbed rotational structure (i.e. Δm)
- $\text{Abs}_{\text{s-pol}} - \text{Abs}_{\text{p-pol}} = \frac{1}{2} \mu_{\perp}^2 - \frac{1}{2} \mu_{\parallel}^2$


Polarization Spectroscopy




Polarization analysis provides:

- An assignment of the perpendicular and parallel components to the crystal field perturbed rotational structure (*i.e.* Δm)
- $\text{Abs}_{\text{s-pol}} - \text{Abs}_{\text{p-pol}} = \frac{1}{2} \mu_{\perp}^2 - \frac{1}{2} \mu_{\parallel}^2$

Rotation in parahydrogen



Simulation:
Linear molecule in *h.c.p* crystal field




Simulation uses theory of: Devonshire, Proc. Roy. Soc. London A 153, 601 (1936);


Bowers and Flygare, J. Chem. Phys., 44, 1389 (1966).

Approved for Public Release; Distribution Unlimited. 96 ABW/PA 06-05-08-287



HCN in parahydrogen




* $o\text{-H}_2$ clustering



HCN/DCN Energy Level Diagrams



Interpretation of the Rotational Fine Structure- HCN + DCN



Crystal field theory (linear molecule in HCP lattice)*

$$H = H_{\text{r,v}} + V_{\text{cry}}$$

$$V_{\text{cry}} = \boxed{\varepsilon_2} C_{2,0}(\Omega_{\text{HCN}}) + \boxed{\varepsilon_3} [C_{3,-3}(\Omega_{\text{HCN}}) - C_{3,3}(\Omega_{\text{HCN}})]$$

$$\text{where } C_{l,m}(\Omega) = \left(\frac{4\pi}{2l+1} \right)^{\frac{1}{2}} Y_{l,m}(\Omega)$$

“Fitting” Results in cm^{-1} : (10 levels, 9 parameters)



	HCN		DCN	
	$p\text{-H}_2$	Gas	$p\text{-H}_2$	Gas
B	1.475(5)	1.478	1.2791(5)	1.207
ΔB	-0.005(5)	-0.010	-0.032(5)	-0.010
D	0.0675(5)	2.910×10^{-6}	0.0654(5)	1.927×10^{-6}
ΔD	0.0032(5)	0.025×10^{-6}	-0.0028(5)	0.017×10^{-6}
v_0	3302.989(5)	3311.4770	2625.093(10)	2630.3033
ε_2	-1.274(5)	-	-2.142(5)	-
$\Delta \varepsilon_2$	-0.193(5)	-	-0.047(5)	-
ε_3	6.850(5)	-	7.500(5)	-
$\Delta \varepsilon_3$	0.767(5)	-	0.028(5)	-

very large! $\times 20,000!$ Identical!


NOTE: Above perturbation does not converge for large D !
Treatment was modified to incorporate centrifugal distortion after crystal field calculation to avoid this problem.



NO in Solid Parahydrogen



Preliminary C₂H₂ in parahydrogen






$E_{R(0)} - E_{P(1)}$ (4B)		<i>pH</i> ₂		
Gas		<i>pH</i> ₂		
Wavenumber	Relative	Wavenumber	Relative	
HCCH	4.71 cm ⁻¹	1.00	3.35 cm ⁻¹	1
DCCH	3.97 cm ⁻¹	0.84	2.74 cm ⁻¹	0.82
DCCD	3.39 cm ⁻¹	0.72	2.47 cm ⁻¹	0.74

- Acetylene IR spectrum ‘ages’... timescale of 30 minutes. Aging appears to be acetylene-acetylene clustering.
- “Rotational spectrum” disappears fairly quickly (but not with annealing!)

Y.P. Lee and collaborators *Chem. Phys. Lett.* **435** 247-251 (2007) – no rotation at ‘high’ [o-H₂]


Preliminary C₂H₂ in parahydrogen




- HCCH Van der Waal diameter ~ 6.4 Å
- *p*-H₂ intermolecular spacing – 3.8 Å
- Substantial overlap of wavefunctions

Prediction of Crystal Field Parameters: Truncation


Pair Potential



+
 $\text{H}_2 - \text{h.c.p. lattice}$



Rotational PES dopant in *h.c.p* H_2 lattice (Rigid)



Crystal field theory (linear molecule in HCP lattice) *

$$H = H_{\text{r,v}} + V_{\text{cry}}$$


$$V_{\text{cry}} = \sum_{l,m}^{\text{m} \leq l} \epsilon_{l,\pm m} \left(\frac{4\pi}{2l+1} \right)^{1/2} Y_{l,\pm m}(\Omega)$$

* SAPT/MC⁺BS, P. Jankowski and K. Szalwicz, *J. Chem. Phys.* **108**, 3554 (1998).


Approved for Public Release; Distribution Unlimited. 96 ABW/PA 06-05-08-287

Prediction of Crystal Field Parameters: Truncation - CO


Rotational PES for CO



$$V_{\text{cry}} = \varepsilon_2 C_{2,0}(\Omega) + \varepsilon_3 [C_{3,-3}(\Omega) - C_{3,3}(\Omega)]$$



CFT –up to $C_{12,12}(\Omega)$






(2.52)*

ε (cm^{-1})	Term	ε (cm^{-1})	Term
4.86	$C_{3,3}(\Omega)$	0.65	$C_{9,3}(\Omega)$
6.74	$C_{5,3}(\Omega)$	0.55	$C_{12,6}(\Omega)$
6.54	$C_{4,0}(\Omega)$	0.46	$C_{8,0}(\Omega)$
3.68	$C_{6,6}(\Omega)$	0.43	$C_{12,12}(\Omega)$
-5.80	$C_{6,0}(\Omega)$	-0.02	$C_{12,0}(\Omega)$
2.61	$C_{7,3}(\Omega)$	-0.47	$C_{11,3}(\Omega)$
1.10	$C_{8,6}(\Omega)$	-0.22	$C_{2,0}(\Omega)$

* Experimentally fitted values.

(1.03)*

Prediction of Crystal Field Parameters: Truncation - HCN




ε (cm^{-1})	Term	ε (cm^{-1})	Term
-3.04	$C_{6,6}(\Omega)$	0.63	$C_{8,6}(\Omega)$
-2.28	$C_{5,3}(\Omega)$	0.09	$C_{8,0}(\Omega)$
3.35	$C_{4,0}(\Omega)$	-0.01	$C_{9,3}(\Omega)$
-4.82	$C_{6,0}(\Omega)$	(6.85/7.50)*	
0.70	$C_{3,3}(\Omega)$	(-1.27/-2.14)*	
-1.24	$C_{7,3}(\Omega)$		
-0.47	$C_{2,0}(\Omega)$		

* Experimental fitted values for HCN/DCN.

* HCN-H₂ Pair Potential - D. Moore, unpublished,

Approved for Public Release; Distribution Unlimited. 96 ABW/PA 06-05-08-287

Estimation of Crystal Field Parameters: Simulated Spectrum for CO + HCN



- Qualitative agreement with experiment
- Overestimates splitting in $J=1$ manifold – rigid lattice structure
- Underestimates $\epsilon_{2,0}$ parameter
- Compression of rotational fine structure (B_{eff}/D_{eff}) not accounted for by crystal field

Summary

Conclusions

- Parahydrogen is remarkably accommodating to rotating molecules... even HCCH!
- “reduction” in rotational constant can be thought as a large centrifugal distortion constant... B may not be affected at all
- Simulations using pair potentials and rigid lattice agree qualitatively with experiment

Acknowledgements:

- Takamasa Momose (Crystal field theory for CO)
- David Anderson (NO data)
- David Moore (H_2 -HCN potential calculations)



QUESTIONS?

